arXiv:1207.1466v2 [cond-mat.quant-gas] 4 Aug 2012

# Vortices on demand in multicomponent Bose-Einstein condensates

R. Zamora-Zamora, M. Lozada-Hidalgo, S. F. Caballero-Benítez and V. Romero-Rochín*
*Instituto de Física, Universidad Nacional Autónoma de México.*
*Apartado Postal 20-364, 01000 México D. F. Mexico.*
(Dated: August 7, 2012)

We present a novel and simple mechanism to produce vortices at any desired spatial locations in harmonically trapped Bose-Einstein condensates (BEC) with multicomponent spin states coupled to external transverse magnetic fields. The vortices appear at the spatial points where the spin-field interaction vanishes and, depending on the multipolar magnetic field order, the vortices acquire a corresponding charge. We explicitly demonstrate our findings, both numerically and analitically, by analyzing 2- and 3D BEC via the Gross-Pitaevskii equation for atomic systems with either two or three internal states. We further show that, by an spontaneous symmetry breaking mechanism, vortices can appear in any spin component, unless symmetry is externally broken at the outset. We propose an experimental realization of this scenario using an ultracold gas of $^{87}$Rb occupying all three $F = 1$ states in an optical trap. We suggest that a further time varying of the external magnetic field may lead to a route for quantum turbulence.

PACS numbers: 67.85.Fg, 03.75.Lm,05.30.Jp

## I. INTRODUCTION

Quantum vortices are the hallmark of superfuidity in either fermionic or bosonic ultracold gases[1–7]. Following the seminal work of Abrikosov[8] and other works on vortices in rotating Helium [9, 10], it has been widely verified that rotating Bose-Einstein condensates (BEC), under appropriate angular velocities of rotation[5–7, 11], indeed show a triangular vortex lattice as the most stable state. Alternative ways have been devised to create vortices, such as phase imprinting[12–15] and, very recently, by creation of artificial gauge fields[16, 17]. In this article we introduce another way of creating vortices; one that necessarily requires the presence of more than one atomic internal spin state coupled to an external transverse magnetic field. In this regard it is of relevance to recall that, even though very early in the development of optical traps[18, 19] it was pointed out[20–22] the additional richness of spinor or multicomponent Bose-Einstein condensates, most of the efforts in creating and studying vortices were focused on rotating condensates[23]. The novelty of our proposal, in addition to exploit the spinorial structure of the condensates, is that the vortices can be created essentially at any desired spatial location. This property, as we shall see, depends on the ability to produce a corresponding on-demand tailored magnetic field.

The essence of vortex appearance follows from the same argument provided by Abrikosov[8]. In superconductors, vortices appear in the spatial points where the type II superconductor allows the magnetic field to penetrate. In those vortices (or flux tubes as they are also known) the density is zero and the superconductor current circulates around the vortex to cancel the field in the bulk. In our proposal the situation is very similar, except that we work with multi-component wavefunctions and, unlike Abrikosov, we work with non-uniform magnetic fields.

We find, interestingly, that the vortices are nucleated not where the magnetic field is able to penetrate the condensate, but on the spatial points where the field is zero. Additionally, the vortex solution will not have the same charge in all spin components of the wave function. That is, if a vortex of charge $Q$ is nucleated on one spin component, the vortex solution on the other spin components for the same spatial location must have a charge $q \neq Q$. Moreover, if this charge is zero, $q = 0$, no vortex will appear and the spin component will develop a density accumulation or spike so as to cancel the vortex formed on the other component(s), leaving the total density unaltered in the usual gaussian-like form. In this way, vortices will be nucleated "on demand" at the spatial positions where the magnetic field is zero, albeit not with the same charge on all spin components.

This approach presents some additional interesting consequences. The first one is that the vortices can be created with arbitrary integer charge, $Q = 1, 2, 3\ldots$, by selecting a field with appropriate multipolar order; dipolar fields give rise to $Q = 1$, quadrupolar ones to $Q = 2$, et cetera. In this way, not only the position of the vortices, but their charge as well, can be controlled using on-demand tailored magnetic fields.

The second interesting consequence of our approach is that it is possible to assemble vortex lattices "on demand" by selecting magnetic fields with zeroes on the lattice sites. Now, since all solutions obtained from the Gross-Pitaevskii and Ginzburg-Landau equations are of minimum free energy, we conclude that triangular vortex lattices observed both at superconductors [24] and BEC alike [5–7, 11], are a peculiarity consequence of the use of *uniform* magnetic fields. Rotating BEC are not an exception to this. It is possible to see that a rotating BEC is equivalent to a superfluid in a *uniform* magnetic field. In this way, our approach may have the further advantage

---

* romero@fisica.unam.mx

of providing evidence of a unique cause underlying the different vortex nucleation techniques currently at use.

In this article we first analyze the corresponding Gross-Pitaevskii (GP) equation to describe how the vortices appear. We consider a very general case and then illustrate several different cases in 2 and 3D gases by numerically solving the corresponding GP equation. In the last section we suggest how this situation can be experimentally realized in a gas of $^{87}$Rb occupying all three $F = 1$ states in an optical trap. In this regard, we point out that precursors of the present study are the analysis of Refs. [25] and [26], the former discussing the appearance of vortices in the center of a Ioffe-Pritchard trap in a $F = 1$ condensate, while the latter being a detailed study of vortices on two-level condensates, yet, in rotating systems.

## II. A MULTICOMPONENT BEC IN A *TRANSVERSAL* MAGNETIC FIELD

Consider a 2 or 3 dimensional BEC superfluid at zero temperature. The gas is confined by an optical trap. In addition, there exists an external *transveral* magnetic field that couples to the $x$ and $y$ spin components. The atoms have spin $J$. The $2J + 1$ GP equations are,

$$\left[-\frac{\hbar^2}{2m}\nabla^2 + V^{\alpha}_{ext}(\vec{r}) + \sum_{\beta} g^{\alpha\beta}\psi^*_{\beta}(\vec{r})\psi_{\beta}(\vec{r})\right]\psi_{\alpha}(\vec{r}) \qquad (1)$$
$$-m_0\sum_{\gamma}\left[J^{\alpha\gamma}_x B_x(\vec{r}) + J^{\alpha\gamma}_y B_y(\vec{r}) + J^{\alpha\gamma}_z B_z\right]\psi_{\gamma}(\vec{r}) = \mu\psi_{\alpha}(\vec{r}),$$

where $\vec{r}$ is the $D$-dimensional position vector. The Greek letters, $\alpha, \beta = 1, 2, \ldots, 2J + 1$, denote spin components. The confining external potentials are $V^{\alpha}_{ext}(\vec{r})$ and we have considered the possibility that each spin component feels a different confining trapping potential; however, for illustration purposes we shall assume below that they are all the same harmonic potential. In general, we assume the external trapping potentials to be smooth at any point. We can also, in principle, assume that the scattering length is different for any pair of components; but again, in the numerical evaluations below, we will consider the simple case $g^{\alpha\beta} = g\delta_{\alpha\beta}$. This is certainly an important point since it is known[27] that the scattering length truly depends on the internal atomic states and this in turn leads to different types of macroscopic "magnetic" properties of the condensates. By assuming a unique positive scattering length throughout, one may assert that the condensates with which we are dealing are at the border of being ferromagnetic or polar[20]. This does not invalidate the point we are highlighting here.

The most important part, and the novelty of our argument, is the coupling of the spin components via an external *inhomogeneous* transversal magnetic field $\vec{B}(\vec{r}) = B_x(\vec{r})\hat{e}_x + B_y(\vec{r})\hat{e}_y$. Accordingly, $J^{\alpha\beta}_x$ and $J^{\alpha\beta}_y$ are the $x$ and $y$ angular momentum matrices of spin $J$. In addition, we also include a *uniform* longitudinal magnetic field $B_z$. We consider here the cases $J = 1/2$ and $J = 1$ only, but the argument may be extended to higher orders. In (2), $\mu$ is the chemical potential and $m_0$ the magnetic moment of the atom.

Now, the external magnetic field must obey $\nabla \cdot \vec{B} = 0$ at any point. In addition, since we want to resemble a true experimental situation, we should demand that throughout the condensate $\nabla \times \vec{B} = 0$ is also obeyed. In what follows we will consider these two restrictions to build the proposed fields, however, we will also include more general situations. We also defer to the following sections the question of how the proposed fields may be produced.

The main point of the discussion is based on proposing that the transversal magnetic field becomes zero at isolated points $\vec{r}_0$, namely, $B_x(\vec{r}_0) = B_y(\vec{r}_0) = 0$. Placing the origin at $\vec{r}_0$, the magnetic field very near that point is assumed to have the following form,

$$B_x(x, y) \approx \mathcal{B}_l\left(Y^{-l}_l(\pi/2, \phi) + (-1)^l Y^l_l(\pi/2, \phi)\right) r^l \qquad (2)$$

and

$$B_y(x, y) \approx -i\mathcal{B}_l\left(Y^{-l}_l(\pi/2, \phi) - (-1)^l Y^l_l(\pi/2, \phi)\right) r^l \qquad (3)$$

where $Y^l_l(\theta, \phi)$ are the usual Spherical Harmonics of order $l = 1, 2, 3\ldots$, $\tan\phi = y/x$ and $r^2 = x^2 + y^2$. Because the fields are constant in the $z$ direction, the fields are to be evaluated at $\theta = \pi/2$. We shall call $l = 1$ dipolar, $l = 2$ quadrupolar, etc.[28]. $\mathcal{B}_l$ are constants with appropriate units. We do not write the constant $z$ component of the field. For the sake of exemplifying the two simplest cases, we write down the explicit forms for $l = 1$ and $l = 2$,

$$\vec{B}(x, y) \approx \mathcal{B}_1\left(x\hat{e}_x - y\hat{e}_y\right) \quad \text{for} \quad l = 1 \qquad (4)$$

and

$$\vec{B}(x, y) \approx \mathcal{B}_2\left((x^2 - y^2)\hat{e}_x - 2xy\hat{e}_y\right) \quad \text{for} \quad l = 2. \qquad (5)$$

Let us turn now to the main problem: vortex nucleation using the fields already discussed. For simplicity we analyze the case of spin $J = 1/2$. This case may be realized with the coupling of the magnetic field to only two components of a larger actual bosonic state, $F > J$ with $F$ integer. In this case, $J_x$ and $J_y$ are the $x$ and $y$ Pauli matrices.

Let us analyze equation (2) in the vicinity of $\vec{r}_0$. In a spinor notation, we denote $\alpha = 1$ as "+" and $\alpha = 2$ as "−". Hence, (2) reduces to the following coupled differential equations,

$$\left[-\frac{\hbar^2}{2m}\nabla^2 + V^{\pm}_{ext}(\vec{r} + \vec{r}_0) + g\left(|\psi_+(\vec{r})|^2 + |\psi_-(\vec{r})|^2\right)\right]\psi_{\pm}(\vec{r})$$
$$+m_0\mathcal{B}_l r^l e^{\pm il\phi}\psi_{\mp}(\vec{r}) = \mu_{\pm}\psi_{\pm}(\vec{r}) \qquad (6)$$

where $\mu_{\pm} = \mu \pm m_0 B_z$. Since the external potential is smooth, we can set $V^{\pm}_{ext}(\vec{r} + \vec{r}_0) \approx V^{\pm}_{ext}(\vec{r}_0)$, namely a constant near $\vec{r}_0$. In such a vicinity, we can fairly see that there exists a solution with cylindrical symmetry of the form,

$$\psi_+(\vec{r}) \approx f_+^l(r)e^{il\phi} \quad \text{and} \quad \psi_-(\vec{r}) \approx f_-^l(r) \tag{7}$$

or, alternatively, with the "+" solution without a phase and with the "−" with the phase $\exp(-il\phi)$.

Substituting the above forms into Eqs.(6) yields two coupled equations for the functions $f_+^l(r)$ and $f_-^l(r)$. The ensuing equations can have a solution only if near $r = 0$ they behave as,

$$f_+^l(r) \approx C_l^+ r^l \quad \text{and} \quad f_-^l(r) \approx C_l^- \quad \text{as} \quad r \to 0. \tag{8}$$

This can be seen by proposing a formal power expansion near $r = 0$ of both functions[8].

The component $\psi_+(\vec{r})$ represents a vortex solution of order (or charge) $l$. This is readily seen by calculating its circulation

$$\mathcal{C}_+ = \oint \vec{v}_s \cdot d\vec{r} = \frac{\hbar}{m} 2\pi l, \tag{9}$$

where $\vec{v}_s$ is the superfluid velocity field and the integration is on any closed contour around $\vec{r}_0$.

The component $\psi_-(\vec{r})$ is not a vortex solution; namely, its circulation vanishes $\mathcal{C}_- = 0$. However, it can be shown that its value at the vortex location $\vec{r}_0$ obeys,

$$|\psi_-(\vec{r}_0)|^2 = \frac{1}{g^{--}} \left(\mu - V_{ext}^-(\vec{r}_0)\right) \tag{10}$$

which is the Thomas-Fermi solution at that point. This is the density accumulation or spike referred to at the introduction.

Had we chosen the solution with the phase in the "−" component, a vortex would have been obtained in that component with negative circulation, $\mathcal{C}_- = -(\hbar/m)2\pi l$. Analogously, $\mathcal{C}_+ = 0$ and no vortex would have been formed for the "+" component.

Within the previous discussion, it is important to note that there is no a priori argument to decide where the vortex will be formed. A BEC with the same potentials, $V^+ = V^-$ can nucleate vortices in either component depending on the initial conditions. There must be a spontaneous symmetry breaking mechanism making the decision. As we shall see below, the constant longitudinal field $B_z$ breaks such a symmetry and the vortices do appear in the positive direction of the field.

Now, let us note that since the above argument only gives the solution at the given point $\vec{r}_0$ where the external magnetic field vanishes, one can consider cases where a magnetic field has zeroes at different spatial locations. Applying the same argument near any of those points, a vortex will be nucleated at any desired position. It is also a simple calculation to verify that the "size" $\xi$ of the vortex is the same one as that of the classical solution of Gross[29] and Pitaesvkii[30], namely $\xi = (\hbar^2/2m\mu)^{1/2}$.

The extension to $J = 1$ is immediate. To see this, consider the magnetic field term only. It can be written as,

$$\sum_\gamma \left[J_x^{\alpha\gamma} B_x(\vec{r}) + J_y^{\alpha\gamma} B_y(\vec{r})\right] \psi_\gamma \approx$$

$$\mathcal{B}_l r^l \begin{pmatrix} 0 & e^{il\phi} & 0 \\ e^{-il\phi} & 0 & e^{il\phi} \\ 0 & e^{-il\phi} & 0 \end{pmatrix} \begin{pmatrix} \psi_{+1} \\ \psi_0 \\ \psi_{-1} \end{pmatrix} \tag{11}$$

This yields a $+l$ vortex at $M_F = +1$, no vortex at $M_F = 0$ and a $-l$ vortex at $M_F = -1$. The opposite solution is also equally valid.

At this point it is relevant to comment on other vortex-like solutions to the GP equation. Because of the non-linearity of the GP equation, other solutions besides the one obtained by variable separation may be valid. Until now we have considered only one kind of solution in both the $J = 1/2$ and $J = 1$ cases. In the former, all the topological charge is concentrated in only one of the spin components "+" or "-"; in the later, equal and opposite charge is found in the "+" and "-" spin components. However, these are not the only valid solutions to the GP equation presently considered. Let us examine the $J = 1/2$ and $J = 1$ cases.

In the $J = 1/2$ case, close to the vortex where the solution behaves as in eq.(7), it can be seen by inspection that the GP equation accepts solutions of the form

$$\psi_+(\vec{r}) \approx f_+^l(r)e^{ia\phi} \quad \text{and} \quad \psi_-(\vec{r}) \approx f_-^l(r)e^{ib\phi} \tag{12}$$

where a and b are integers and $a = b + l$. In this way, there are $l + 1$ solutions to the GP equation presently considered.

In the $J = 1$ case, the rule to select possible solutions is slightly different. Using similar arguments to those leading to eq.(12), it can be shown that solutions of the form

$$\psi_+(\vec{r}) \approx f_+^l(r)e^{ia\phi}, \psi_0(\vec{r}) \approx f_0^l(r)e^{ic\phi}, \psi_-(\vec{r}) \approx f_-^l(r)e^{ib\phi} \tag{13}$$

where a,b and c are integers, $a = c + l$ and $b = c - l$, are also valid solutions. In this way, in the $J = 1$ case under an "$l$" multipole field, there are $l + 1$ solutions as well.

## III. A GALLERY OF VORTICES ON DEMAND

In this section we illustrate the present mechanism for vortex formation with a variety of cases in both 2D and 3D; additional details case by case are also provided. For simplicity, several assumptions are made. The trapping potential is assumed to be an isotropic harmonic potential for all spin components with the same frequency $\omega$. In all cases the same scattering length is assumed for all collisions, namely, $g^{\alpha\beta} = g\delta_{\alpha\beta}$, with $g$ a free parameter. Units in which $\hbar = m = \omega = 1$ are used and $\kappa_l = m_0\mathcal{B}_l$ is defined as the parameter that determines the strength of the external magnetic field. Throughout, the field in the $z$ component is assumed constant, $B_z = \mathcal{B}_l z_0$. In all cases below, the corresponding set of GP-equations is numerically solved with the method proposed by Zeng and Zhang[31]. The chemical potential is part of the solution and it is also our convergence parameter; its value with its corresponding error is given in all cases shown. The density plots as well as phase-velocity fields $\vec{v}_s = \nabla\Phi$, with $\Phi$ the corresponding phase of the $\psi_\alpha(\vec{r})$ component, are also shown; the circulation is plotted as well.

### A. Single $l = 1$ and $l = 2$ vortices in 2 and 3D BEC

This is the simplest case to consider and we begin with two internal states, namely $J = 1/2$. Letting $l = 1$ in Eqs.(2) and (3), the field is dipolar in the sense that it is a linear combination of spherical harmonics of order one. It is very similar to the Ioffe-Pritchard trap field except that our field has a constant $z$ component and, consequently, the sign of the y component must change to preserve Maxwell's equations.

Fig. 1 shows a vortex at the origin $\vec{r}_0 = 0$ in a 2D BEC and Fig. 2 a vortex at an arbitrary $\vec{r}_0$, see figure caption for details. As mentioned above, the density and phase-velocity plots, as well as numerical calculation of circulations, are also shown. Square contours around the vortex were used for numerical integration of the circulation; as the contour includes enough points, one of the circulations gives $\mathcal{C}/2\pi = \pm 1$ while the other gives zero within numerical accuracy.

The population of both states, $\rho_\pm = |\psi_\pm(\vec{r})|^2$ and the total one, $\rho_+ + \rho_-$, are also indicated in the figures. The values for both the "+" and "−" states are close to 0.5 but not exactly. We have no explanation for these deviations from half-and-half population. It can also be noted from the figures that if a vortex is nucleated on one of the spin components, no vortex will be nucleated on the same spatial position in the other components. Additionally, the other spin components develop a density accumulation or spike that cancel the vortex formed in the other component, leaving the total wave function in the usual gaussian-like solution.

Note that in Figs. 1 and 2 the vortices appear in the "−" component. We have verified that, depending in the arbitrary initial conditions of the numerical method, the vortices may appear in any of them. As mentioned above, we believe there is an spontaneous symmetry-breaking mechanism acting here. However, as shown in Fig. 3, if the *longitudinal* magnetic field $B_z = \mathcal{B}_1 z_0$ is different from zero, for instance $z_0 = \pm 5$, the vortices always appear in the "±" component respectively.

Fig. 4 is the 3D version of Fig. 1. We find that although the clouds have spherical symmetry, inherited from the trapping potential, the vortices are of cylindrical nature. This is because the nucleation points (where the magnetic field becomes zero) are actually filaments that cross the full BEC cloud.

Fig. 5 shows single $l = 2$ vortices in a 2D BEC. To produce them, let $l = 2$ in Eqs.(2) and (3). Again, the vortices appear at the chosen vanishing points of the magnetic field. However, the vortices have charge $l = 2$, namely, the circulation gives $\mathcal{C}/2\pi = \pm 2$.



FIG. 1. Vortex of charge $l = 1$ at the origin $\vec{r}_0 = 0$ with $J = 1/2$. We show density plots, (a) "+" component, (b) "-" component, and (c) total density; circulation fields, (d) "+" component, and (e) "-" component; total circulation values as function of the radius of integrations circuit. Parameters are: $g = 4000$, $\kappa = 1.0$ and $z_0 = 0$ (i.e. no longitudinal field). The chemical potencial is $\mu = 29.9594 \pm 0.0001$, and the total relative spin populations are $\frac{<N_+>}{N} = 0.51$ and $\frac{<N_->}{N} = 0.49$. The vortex is in "−" component and has circulation $\mathcal{C}/2\pi = -1$.

Results for $J = 1$ are reported in Fig. 6. The simplest case corresponds to a single $l = 1$ vortex at the origin. As predicted in Section II, the $\alpha = 1$ and $\alpha = -1$ components show single vortices with opposite sign circulations of charge 1; the $\alpha = 0$ component does not present a vortex. In this case, our method does not give rise to other charge distributions within the spin components; however, some may be found in the literature [25].



FIG. 2. Vortex of charge $l = 1$ at the chosen place $\vec{r}_0 = (4, 2)$, with $J = 1/2$. We show density plots, (a) "+" component, (b) "-" component, and (c) total density; circulation fields, (d) "+" component, and (e) '-" component; total circulation values as function of the radius of integrations circuit. Parameters are: $g = 8000$, $\kappa = 0.5$ and $z_0 = 0$. The chemical potencial is $\mu = 46.5984 \pm 0.0002$, and the relative spin population are $\frac{<N_+>}{N} = 0.51$ and $\frac{<N_->}{N} = 0.49$. (b) The vortex is in "−" component and has circulation $\mathcal{C}/2\pi = -1$.

### B. Multiple $l = 1$ vortices in 2D BEC

It was reported in the previous subsection how external fields of the form given by Eqs.(2) and (3) nucleate single vortices at the desired locations. We now turn to the possibility of nucleating arbitrary arrays of vortices.

The first case is shown in Fig. 7, where we study a $J = 1/2$ 2D BEC. The external magnetic field is a superposition of fields generated by eight "infinite" wires (with currents of different magnitudes and signs) placed *outside* the BEC cloud such that $\nabla \cdot \vec{B} = 0$ and $\nabla \times \vec{B} = 0$ inside the cloud region. A single wire located at $\vec{r}_0$ has a field,

$$\vec{B}(x, y) = I \frac{(x - x_0)\hat{e}_y - (y - y_0)\hat{e}_x}{(x - x_0)^2 + (y - y_0)^2} \tag{14}$$

where $I$ is the current in dimensionless units, and $\hat{e}_\xi$ is the unitary vector in direction $\xi$.

With the appropriate choice of the currents $I$ (see the figure caption for details), we were able to generate three spatial points where the field vanishes linearly, namely



FIG. 3. Vortices of charge $l = 1$ at the origin $\vec{r}_0 = 0$, with $J = 1/2$. (a,b) $z_0 = 1$, the vortex appears in the + component, with positive charge +1; chemical potencial is $\mu = 29.8472 \pm 0.0001$ and the populations are $\frac{<N_+>}{N} = 0.37$ and $\frac{<N_->}{N} = 0.63$. (c,d) $z_0 = -1$, the location, population and circulation of the vortex is reversed; chemical potencial is $\mu = 29.8562 \pm 0.0006$, populations are $\frac{<N_+>}{N} = 0.63$ and $\frac{<N_->}{N} = 0.37$



FIG. 4. Vortex of charge $l = 1$ at the origin $\vec{r}_0 = 0$, with $J = 1/2$ and $z_0 = 0$, for a 3D BEC. The vortex is in the "−" component with negative circulation. Chemical potencial is $\mu = 15.48 \pm 0.01$; populations are $\frac{<N_+>}{N} = 0.53$ and $\frac{<N_->}{N} = 0.47$.



FIG. 5. Vortex of charge $l = 2$ at the origin $\vec{r}_0 = 0$, with $J = 1/2$, $z_0 = 0$, $g = 1000$ and $\kappa = 0.2$. The vortex is in the "−" component with negative circulation $\mathcal{C}/2\pi = -2$. Chemical potential is $\mu = 13.9 \pm 0.3$. Populations are $\frac{<N_+>}{N} = 0.58$ and $\frac{<N_->}{N} = 0.4$.



FIG. 6. Vortices of charge $l = 1$ at the origin $\vec{r}_0 = 0$, with $J = 1$, $z_0 = 0$, $g = 8000$ and $\kappa = 0.5$. Chemical potential is $\mu = 47.1020 \pm .0003$; populations are $\frac{<N_+>}{N} = 0.25$, $\frac{<N_->}{N} = 0.25$ and $\frac{<N_0>}{N} = 0.50$ . The vortices appear at the $m_J = \pm 1$ (a,b,d,e). There is no vortex at $m_J = 0$ (c). The vortex at $m_J = +1$ has positive circulation $\mathcal{C}/2\pi = +1$, while the circulation is negative for $m_J = -1$ (f) .

$l = 1$ "zeroes". This should generate three vortices at those locations and, as shown in Fig. 7, we do find 3 vortices with $l = \pm 1$ charge. For the case $z_0 = 0$, two vortices appear in one component and one in the other. Nonetheless, in some cases the three vortices appear in only one component. Again, we have no explanation at this moment for these occurrences and we appeal once more to a symmetry breaking mechanism. However, if $z_0 \neq 0$, the three vortices appear (not shown) at the component corresponding to the sign of $z_0$.

Fig. 8 shows a regular lattice of vortices. These were generated by a magnetic field of the form,

$$\vec{B}(x, y) = \mathcal{B}\left(\sin(\pi x/\lambda)\cos(\pi y/\lambda)\hat{e}_x - \sin(\pi y/\lambda)\cos(\pi x/\lambda)\hat{e}_y\right). \tag{15}$$

The field vanishes linearly at the points $(x, y) = (\pm n\lambda, \pm m\lambda)$ with $n$ and $m$ both integers or both half integers; that is, those points are "zeroes" of dipolar order $l = 1$.

Another interesting feature of this lattice is that vortices of charge $l = \pm 1$ are generated within the same spin component. In the zeroes of the magnetic field that correspond to both $m$ and $n$ integers, vortices of charge $l = +1$ are nucleated in the "+" spin component and with charge $l = -1$ in the "−" spin component. In the same fashion, in those zeroes corresponding to $m$ and $n$ both half integers, vortices of charge $l = -1$ are nucleated in the "+" spin component and with charge $l = +1$ in the "−" spin component. Fig. 8 illustrates a vortex lattice of charge $l = \pm 1$ formed in the "+" spin component, following the rules just described.

Such a behavior can be readily understood using the tools from Section II. Expanding the magnetic field around each of its zeroes, we note that the behavior of



FIG. 7. Three vortices of charge $l = 1$ at the chosen locations $\vec{r}_1 = (3.6, -6.1)$, $\vec{r}_2 = (-6.6, 0)$ and $\vec{r}_3 = (3.6, 6.1)$, with $J = 1/2$, $z_0 = 0$ and $\kappa = 2.0$. We show density plots, (a) "+" component, (b) "-" component, and (c) total density; circulation fields, (d) "+" component, and (e) '-" component; total circulation values as function of the radius of integrations circuit. Note that the vortex in the "+" component has positive +1 circulation, while those in the "−" component each has -1 circulation. Chemical potencial is $\mu = 53.4084 \pm 0.0002$; population are $\frac{<N_+>}{N} = 0.525$ and $\frac{<N_->}{N} = 0.475$.

the magnetic term in Eq.(2) depends on the nature of the zeores. The magnetic term is

$$A^{\pm}_{mn} m_0 \mathcal{B} r \begin{pmatrix} 0 & e^{\pm i\phi} \\ e^{\mp i\phi} & 0 \end{pmatrix} \tag{16}$$

where $A^{\pm}_{mn}$ is a constant. The first case corresponds to both $n$ and $m$ integers, while the second one to the half integer case. Again, the lattice may appear part in one spin component and part in another, unless a longitudinal magnetic field is applied. An "undesired" characteristic of this lattice is that $\nabla \times \vec{B}$ is not zero in the whole region occupied by the BEC cloud. That is, one needs "sources" in the region occupied by the condensate. Nevertheless, this case illustrates the possibility of generating vortices at the desired locations. We leave the tailoring of the corresponding fields to the ingenuity of the experimental researchers. This case can also be realized in a 3D BEC.



FIG. 8. Vortex lattices of charge $l = 1$ for a $J = 1/2$ BEC, see text for magnetic field. In all cases $g = 8000$ $\kappa = 2$. (a,c) $z_0 = 0$ showing the spontaneous symmetry breaking mechanism with chemical potencial $\mu = 49.5221 \pm 0.0003$ and populations $\frac{<N_+>}{N} = 0.62$ and $\frac{<N_->}{N} = 0.38$. (b,d) $z_0 = +5$ while (c,f) are for $z_0 = -5$ with chemical potencial $\mu = 40.3934 \pm 0.0006$ and $\mu = 40.3937 \pm 0.0006$ respectively. Populations in (b,d) are $\frac{<N_+>}{N} = 0.004$ and $\frac{<N_->}{N} = 0.996$; for (c,f) populations are $\frac{<N_+>}{N} = 0.996$ and $\frac{<N_->}{N} = 0.004$

## IV. FINAL REMARKS

As we have seen, in multicomponent BEC clouds, the presence of transversal magnetic fields with vanishing values at given spatial locations, nucleate vortices at those very points. The vortices can thus be placed at



FIG. 9. Vortex lattice corresponding to (b) in Fig. 8. The circulation field is shown in black arrows. One can observe the alternating character of the vortex topological charge.

any location "on demand". We have verified this result for many tailored magnetic fields. In addition, we have verified the further prediction that the total density $\rho(x,y) = |\psi_+(x,y)|^2 + |\psi_-(x,y)|^2$ does not show any evidence of the vortices and that its form is the "simple" solution of GP in a harmonic potential.

It is interesting to contrast the present form of the appearance of the vortices with that of a rotating superfluid. In the rotating frame, the (one-component) GP equation looks like[2],

$$-\frac{\hbar^2}{2m}\nabla^2\Phi(\vec{r}) + (\vec{\Omega}\cdot\vec{L})\Phi(\vec{r}) + g\,|\Phi(\vec{r})|^2\,\Phi(\vec{r}) - \mu\Phi(\vec{r}) = 0, \tag{17}$$

where $\vec{\Omega} = \Omega\hat{z}$ is the angular velocity of rotation and $\vec{L}$ is the angular momentum operator,

$$\vec{L} = \frac{\hbar}{i}\vec{r}\times\nabla. \tag{18}$$

By a vector identity one can rewrite,

$$(\vec{\Omega}\cdot\vec{L})\Phi(\vec{r}) = \frac{\hbar}{i}\left(\vec{\Omega}\times\vec{r}\right)\cdot\nabla\Phi(\vec{r}) \tag{19}$$

and, therefore, one can see that the rotation term is equivalent to introducing a vector potential $\vec{A} = \vec{\Omega}\times\vec{r}$ corresponding to a longitudinal uniform "magnetic field" $\vec{B} = \vec{\Omega}$ (in appropriate units).

In its celebrated paper, Abrikosov[8] showed that if the above equations hold, a lattice of vortices, of the type described in Section I, will be created by the presence of such a magnetic or appropriate gauge field. It does appear, therefore, that vortices are created by the coupling of the *orbital* angular momentum $\vec{L}$ with the external field $\Omega$, i.e. $\vec{\Omega}\cdot\vec{L}$. The fact that the vortices appear in a lattice is "incidental" due to the global structure of the GP equation in the presence of a *uniform* longitudinal magnetic field.

In the case described in this work, vortices also appear because of a magnetic angular momentum coupled to an external gauge field. It does appear, however, that in the present case it is the *intrinsic* or *spin* angular momentum the main player instead of the orbital angular momentum.

We believe that the above two cases may not be as different as they may appear. This assertion may be based on the observation that GP equation near a zero of the magnetic field can be transformed as one with an artificial gauge field coupled to the orbital angular momentum. Let us see this briefly. For simplicity, take the case $J = 1/2$, $z_0 = 0$ and $g_{\alpha\beta} = g\delta_{\alpha\beta}$. The coupled equations (2) may be written as

$$\left[-\frac{\hbar^2}{2m}\nabla^2 + V_{ext}(\vec{r}) + g|\Psi|^2 + \kappa\left(x\sigma_x + y\sigma_y\right) - \mu\right]\Psi = 0. \tag{20}$$

where ,

$$\Psi = \begin{pmatrix}\psi_+\\ \psi_-\end{pmatrix}, \tag{21}$$

and $|\Psi|^2 = |\psi_+|^2 + |\psi_-|^2$. By applying the unitary transformation

$$U = e^{-i\phi\sigma_z/2}e^{-i\pi\sigma_y/4}e^{i\phi\sigma_z/2}, \tag{22}$$

with $\tan\phi = y/x$, equation (20) now becomes[33, 34]

$$\begin{aligned}\Big[\frac{1}{2m}&(-i\hbar\nabla - \vec{A}_x\sigma_x - \vec{A}_y\sigma_y - \vec{A}_z\sigma_z)^2\\ &+V_{ext}(\vec{r}) + g|\tilde{\Psi}|^2 + \kappa r\sigma_z - \mu\Big]\tilde{\Psi} = 0,\end{aligned} \tag{23}$$

with $\tilde{\Psi}$ the transformed state. The induced artificial gauge potentials may be easily calculated[33]. In particular, the diagonal potential $\vec{A}_z$ is,

$$\vec{A}_z = -\frac{\hbar}{2r}\hat{e}_\phi \tag{24}$$

which corresponds to a nonuniform *singular* magnetic field, actually, to that of a magnetic monopole. Thus, the vortex nucleation in the rotating condensate appears to be equivalent to ours via an artificial magnetic monopole field. This, in turn, is intimately related to Berry phases[36, 45]. Therefore, the present scheme may also be seen from the perspective of artificial gauge fields[16, 17, 32, 46, 51], although it should be clear that we do not assume adiabaticity of the diagonal terms in (23).

We believe the present results can be simply realized in actual experimental conditions. In particular, for a $^{87}$Rb BEC there are two cases that can be simply matched. One is the $F = 1$ case that naturally corresponds to the $J = 1$ situation here discussed.The other, the case $F = 2$ may be tailored to have only the components $m_1 = 1$ and $m_2 = 2$ corresponding to $J = 1/2$. The external magnetic fields may be produced by single-wire magnetic fields, as given by (14). However, an additional nonuniform small magnetic field in the $z$ direction should be used to separate the spin components such that the phenomena may be observed.

The present scheme may also be used as a route to quantum turbulence. Recently, Bagnato et al.[39] found a way to stir a $^{87}$Rb condensate by means of time dependent magnetic fields, achieving a state of quantum turbulence. Although we are not asserting that our procedure explains those experimental results, it does appear that once vortices are nucleated by transversal magnetic fields, the vortices may be stirred by simply changing in time the positions of the zeroes of the magnetic field. It seems evident that a strong stirring of the vortices may lead to a quantum turbulent state. The study is under way and will be reported elsewhere.

We thank support from grant DGAPA UNAM IN108812. R.Z.Z. and M.L.H acknowledge support from CONACYT, Mexico.

[1] R. Dum, J. I. Cirac, M. Lewenstein and P. Zoller. *Phys. Rev. Lett.* **80**, 2972 (1998).
[2] Y. Castin and R. Dum. *Eur. Phys. Journal D* **7**, 399 (1999).
[3] M. R. Matthews et. al. *Phys. Rev. Lett.* **83**, 2498 (1999).
[4] M. W. Zwierlein, J. R. Abo-Shaeer, A. Schirotzek, C. H. Shunck and W. Ketterle. *Nature* **435**, 1047 (2005).
[5] N. R. Cooper. *Adv. Phys.* **57**, 539 (2008).
[6] J. R. Abo-Shaeer, C. Raman, J. M. Vogels and W. Ketterle. *Science* **292**, 476 (2001).
[7] K. W. Madison, F. Chevy, W. Wohlleben and J. Dalibard. *Phys. Rev. Lett.* **84**, 806 (2000).
[8] A. A. Abrikosov. *JETP* **32**, 1442 (1952).
[9] E. J. Yarmchuk, M. J. V. Gordon and R. E. Packard. *Phys. Rev. Lett.* **43**, 214 (1979).
[10] R. J. Donnelly. *Quantized Vortices in Helium II.* Cambridge Studies in Low Temperatrue Physics (Cambridge, 1991).
[11] V. Schweikhard, I. Coddington, P. Engels, V. P. Mogendorff and E. A. Cornell. *Phys. Rev. Lett.* **92**, 040404 (2004).
[12] A. E. Leanhardt, A. Görlitz, A. P. Chikkatur, D. Kielpinski, Y. Shin, D. E. Pritchard and W. Ketterle. *Phys. Rev. Lett.* **89**, 190403 (2002).
[13] M. Nakahara, T. Isoshima, K. Machida, S. Ogawa, and T. Ohmi. *Physica B*, 284-288 (2000) 17.
[14] T. Isoshima, M. Nakahara, T. Ohmi, and K. Machida. *Phys. Rev. A* **61**, 063610 (2000).
[15] K. C. Wright, L. S. Leslie, A. Hansen, N. P. Bigelow. *Phys. Rev. Lett.* **102**, 030405 (2009).
[16] Y. -J. Lin, R. L. Compton, K. Jiménez-García, J. V. Porto and I. B. Spielman. *Nature Letters* **462**, 628 (2009).
[17] K. J. Gunter, M. Cheneau, T. Yefsah, S. P. Rath and J. Dalibard. *Phys. Rev. A* **79**, 011604(R) (2009).
[18] J. Stenger, S. Inouye, D. M. Stamper-Kurn, H.-J. Miesner, A. P. Chikkatur and W. Ketterle. *Nature* **396**, 345 (1998).
[19] H. -J. Miesner, D. M. Stamper-Kurn, J. Stenger, S. Inouye, A. P. Chikkatur, and W. Ketterle. *Phys. Rev. Lett.* **82**, 2228 (1999).
[20] T. -L. Ho. *Phys. Rev. Lett.* **81**, 742 (1998).
[21] T. Ohmi and K. Machida. *J. Phys. Soc. Jap.* **67**, 1822 (1998).
[22] S. -K. Yip. *Phys. Rev. Lett.* **83**, 4677 (1999).
[23] A. L. Fetter. *Rev. Mod. Phys.* **81** 647 (2009)
[24] U. Essmann and H. Trauble. *Phys. Lett. A* **24**, 526 (1967).
[25] E. N. Bulgakov and A. F. Sadreev. *Phys. Rev. Lett.* **90**, 200401 (2003).
[26] K. Kasamatsu, M. Tsubota, and M. Ueda. *Int. J. Mod. Phys. B* **19**, 1835 (2005).
[27] N. N. Klausen, J. L. Bohn, and C. H. Greene. *Phys. Rev. A* **64** 053602 (2001).
[28] W. Greiner. *Classical Electrodynamics*, (Springer,1998).
[29] E. P. Gross. *Il nuovo cimento* **20** (1961).
[30] L. P. Pitaevskii. *JETP* **13** (1961).
[31] R. Zeng and Y. Zhang. *Comp. Phys. Comm.* **180**, 854-860 (2009).
[32] I. B. Spielman. *Phys. Rev. A* **79**, 063613 (2009).
[33] V. Romero-Rochin and J.A. Cina. *J. Chem. Phys.* **91**, 6103 (1989).
[34] H. C. Longuet-Higgins, U. Opik, M. H. L. Pryce and R. A. Sack. *Proc. R. Soc. Lond. A*, 1 (1958).
[35] A. A. Abrikosov. Nobel Lecture, December 8, 2003.
[36] J. Larson and E. Sjöqvist. *Phys. Rev. A* **79**, 043627 (2009).
[37] D. V. Freilich, D. M. Bianchi, A. M. Kauffman, T. K. Languin and D. S. Hall. *Science* **329**, 1182 (2010).
[38] W. Bao, I. Chern and F. Lim. *Journal of Computational Physics* **219**, 836 (2006).
[39] E. A. L. Henn, J. A. Seman, G. Roati, K. M. F. Magalhães, and V. S. Bagnato. *Phys. Rev. Lett.* **103**, 045301 (2009).
[40] C. J. Pethick and H. Smith. Bose-Einstein Condensation in Dilute Gases, Second Edition, (Cambridge University Press, 2008).
[41] M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch and I. Bloch. *Nature* **415**, 39 (2003).
[42] C. A. Regal, M. Greiner and D. S. Jin. *Phys. Rev. Lett.* **92**, 040403 (2004).
[43] D. C. Tsui, H. L. Stormer and A. C. Gossard. *Phys. Rev. Lett.* **48**, 1559 (1982).
[44] D. Jaksch and P. Zoller. *N. J. Phys.* **5**, 56 (2003).
[45] M. V. Berry. *Proc. R. Soc. Lond. A* **392**, 45 (1984).
[46] G. Juzeliunas, J. Ruseckas, P. Öhberg and M. Fleischhauer. *Phys. Rev. A* **73**, 025602 (2006).
[47] M. Cheneau et. al. *Europhys. Lett.* **83**, 60001 (2008).
[48] M. F. Andersen et. al. *Phys. Rev. Lett.* **97**, 170406 (2006).
[49] Y.-J. Lin et.al. *Phys. Rev. Lett.* **102**, 130401 (2009).
[50] E. Lundh, C. J. Pethick and H. Smith. *Phys. Rev. A* **55**, 2126 (1997).
[51] D. R. Murray, P. Öhberg, D. Gomila and S. M. Barnett. *Phys. Rev. A* **79**, 063618 (2009).
[52] Y.-J.Lin, A. R. Perry, R. L. Compton, I. B. Spielman and J. V. Porto. *Phys. Rev. A* **79**, 063631 (2009).
[53] P. B. Blakie, A. S. Bradley, M. J. Davies, R. J. Ballagh and C. W. Gardiner. *Adv. Phys.* **57**, 363 (2008).